PATIENT NAME:	:	ID#:	<u>-</u>	DATE:
	survey is meant to help us obtability. Please circle the ans			nts regarding their current levels of
1. Please rate yc	our pain level with activ	vity: NO PAÍN = 0 1	2 3	4 5 6 7 8 9 10 = VERY SEVERE PAIN
MODIFIED OS	SWESTRY DISABILIT	<u> FY SCALE – INI</u>	<u> </u>	<u>/ISIT</u>
(1) The pain is bad, be pain medication. (2) Pain medication p (3) Pain medication p (4) Pain medication p (5) Pain medication p (6) I can take care of (7) I can take care of (8) I can take care of (9) I can take care of (1) I can take care of (1) I need help, but I (2) I is painful to tal (3) I need help every (5) I do not get dress (9) I can lift heavy w (1) I can lift heavy w (2) Pain prevents medicate but I can manage (6) (9) Pain prevents medicate but I can lift only ve (1) I can lift only ve (2) I cannot lift or can (3) Pain prevents m (4) I can lift only ve (5) I cannot lift or can (6) Pain prevents m (7) Pain prevents m (8) Pain prevents m (9) Pain prevents m (1) Pain prevents m (2) Pain prevents m (3) Pain prevents m (4) I can only walk (5) I am in bed mos (6) I can sit in any of (7) I can only sit in	provides me with complete relief provides me with moderate relief provides me with little relief frohas no effect on my pain.  (washing, dressing, etc.)  If myself normally without causing myself normally, but it increase ake care of myself, and I am slow I am able to manage most of my day in most aspects of my care used, wash with difficulty, and state weights without increased pain.  Weights weights are conveniently the from lifting heavy weights off the weights are conveniently are from lifting heavy weights.  Carry anything at all.  Therefore walking more than 1 milling from walking more than 1/2 milling fro	g to take  ef from pain.  ef from pain.  om pain.  ing increased pain.  ses my pain.  w and careful.  y personal care.  e.  tay in bed.  pain.  f the floor,  y positioned  at I can manage y positioned.  ttance.  ile.  ille.  itle.  to the toilet.	(0) (1) (2) (3) (4) (5) 7. (0) (1) (2) (3) (4) (5) 8. (0) (1) (2) (3) (4) (5) 9. (0) (1) (2) (3) (4) (5)	Pain prevents me from standing more than 10 minutes. Pain prevents me from standing at all.  Sleeping Pain does not prevent me from sleeping well. I can sleep well only by using pain medication. Even when I take pain medication, I sleep less than 6 hours. Even when I take pain medication, I sleep less than 4 hours. Even when I take pain medication, I sleep less than 2 hour Pain prevents me from sleeping at all.  Social Life My social life is normal and does not increase my pain. My social life is normal, but it increases my level of pain. Pain prevents me from participating in more energetic activities (eg. sports, dancing). Pain prevents me from going out very often. Pain has restricted my social life to my home. I have hardly any social life because of my pain.  Traveling I can travel anywhere without increased pain. I can travel anywhere but it increases my pain. My pain restricts my travel over 2 hours. My pain restricts my travel over 1 hour. My pain restricts my travel to short necessary journeys journeys under 1/2 hour. My pain prevents all travel except for visits to the physician/therapist or hospital.  Employment / Homemaking My normal homemaking/job activities do not cause pain. My normal homemaking/job activities increase my pain, but I can still perform all that is required of me.
<ul><li>(3) Pain prevents π</li><li>(4) Pain prevents π</li></ul>	Pain prevents me from sitting more than 1 hour. Pain prevents me from sitting more than ½ hour. Pain prevents me from sitting more than 10 minutes. Pain prevents me from sitting at all.		(3) (4) (5)	<ol> <li>I can perform most of my homemaking/job duties, but pain prevents me from performing more physically stressful activities (eg., lifting, vacuuming).</li> <li>Pain prevents me from doing anything but light duties.</li> <li>Pain prevents me from doing even light duties.</li> <li>Pain prevents me from performing any job or homemaking hores.</li> </ol>
ODI © Jeremy Fa France. E-mail: c	contact@mapi-trust.org - I	Internet: <u>www.mapi-tru</u>	formation ust.org	n and permission to use: MAPI Research Trust, Lyon
Comorbidities:	□Cancer □Diabetes	□Obesity		inson's, Muscular Dystrophy, Huntington's, CVA, Alzheimer's, TBI
	☐ Heart Condition ☐ High Blood Pressure ☐ Multiple Treatment Areas	☐ Surgery for this Problem ☐ Systemic Disorders (e.g.		Rheumatoid Arthritis. Fibromyalgia)